

## Bibliography

---

- Alberch, P., Gould, S. J., Oster, G. F. and Wake, D. B. (1979). Size and shape in ontogeny and phylogeny. *Paleobiology*, **5**, 296–317.
- Albrecht, G. (1978). Some comments on the use of ratios. *Systematic Zoology*, **27**, 67–71.
- Anderson, T. W. (1958). *An Introduction to Multivariate Analysis*. Wiley.
- Anstey, R. L. and Pachut, J. F. (1995). Phylogeny, diversity history, and speciation in Paleozoic Bryozoans. In *New Approaches to Speciation in the Fossil Record* (D. H. Erwin and R. L. Anstey, eds) pp. 239–284. Columbia University Press.
- Archie, J. W. (1985). Methods for coding variable morphological features for numerical taxonomic analysis. *Systematic Zoology*, **34**, 236–345.
- Arnold, S. J. and Phillips, P. C. (1999). Hierarchical comparison of genetic variance–covariance matrices. II. Coastal–island divergence in the garter snake, *Thamnophis elegans*. *Evolution*, **53**, 1516–1527.
- Atchley, W. R. and Anderson, D. (1978). Ratios and the statistical analysis of biological data. *Systematic Zoology*, **27**, 71–78.
- Atchley, W. R., Gaskins, C. T. and Anderson, D. (1976). Statistical properties of ratios. I. Empirical results. *Systematic Zoology*, **25**, 137–148.
- Bartlett, M. S. (1947). Multivariate analysis. *Journal of the Royal Statistical Society, Series B*, **8**, 176–197.
- Bastir, M., Rosas, A. and Sheets, H. D. (2004). The morphological integration of the hominoid skull: a partial least squares and PC analysis with morphogenetic implications for European Mid-Pleistocene mandibles. In *Developments in Primatology: Progress and Prospects* (D. Slice, ed.), in press. Kluwer Academic/Plenum Press.
- Berg, R. L. (1960). The ecological significance of correlation pleiades. *Evolution*, **14**, 171–180.
- Bininda-Emonds, O. R. P., Jeffrey, J. E. and Richardson, M. K. (2003). Inverting the hourglass: quantitative evidence against the phylotypic stage in vertebrate development. *Proceedings of the Royal Society of London, Series B*, **270**, 341–346.
- Bookstein, F. L. (1982). The geometric meaning of soft modeling with some generalizations. In *Systems Under Indirect Observation: Causality–Structure–Prediction* (K. G. Jöreskog and H. Wold, eds) pp. 55–74. North Holland Publishing Company.
- Bookstein, F. L. (1986). Size and shape spaces for landmark data in two dimensions. *Statistical Science*, **1**, 181–242.
- Bookstein, F. L. (1989a). Principal warps: thin-plate splines and the decomposition of deformations. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, **11**, 567–585.
- Bookstein, F. L. (1989b). “Size and shape”: a comment on semantics. *Systematic Zoology*, **38**, 173–190.
- Bookstein, F. L. (1991). *Morphometric Tools for Landmark Data: Geometry and Biology*. Cambridge University Press.
- Bookstein, F. L. (1996a). Combining the tools of geometric morphometrics. In *Advances in Morphometrics* (L. F. Marcus, M. Corti, A. Loy et al., eds) pp. 131–152. Plenum Press.
- Bookstein, F. L. (1996b). Standard formula for the uniform shape component in landmark data. In *Advances in Morphometrics* (L. F. Marcus, M. Corti, A. Loy et al., eds) pp. 153–168. Plenum Press.
- Bookstein, F. L. (1997a). Landmark methods for forms without landmarks: morphometrics of group differences in outline shape. *Medical Image Analysis*, **1**, 225–243.

- Bookstein, F. L. (1997b). Shape and the information in medical images: a decade of the morphometric synthesis. *Computer Vision and Image Understanding*, **66**, 97–118.
- Bookstein, F. L., Chernoff, B., Elder, R. et al. (1985). *Morphometrics in Evolutionary Biology*. The Academy of Natural Sciences of Philadelphia.
- Bookstein, F. L., Gunz, P., Ingeborg, H. et al. (2003). Cranial integration in *Homo*: singular warps analysis of the midsagittal plane in ontogeny and evolution. *Journal of Human Evolution*, **44**, 167–187.
- Campbell, N. A. and Atchley, W. R. (1981). The geometry of canonical variates analysis. *Systematic Zoology*, **30**, 268–280.
- Chapman, R. E. (1990). Conventional Procrustes methods. In *Proceedings of the Michigan Morphometrics Workshop* (F. J. Rohlf and F. L. Bookstein, eds) pp. 251–267. University of Michigan Museum of Zoology.
- Chappill, J. A. (1989). Quantitative characters in phylogenetic analysis. *Cladistics*, **5**, 217–234.
- Chatfield, C. and Collins, A. J. (1980). *Introduction to Multivariate Analysis*. Chapman & Hall.
- Cheverud, J. M. (1982). Phenotypic, genetic and environmental integration in the cranium. *Evolution*, **36**, 499–512.
- Cheverud, J. M. (1984). Quantitative genetics and developmental constraints on evolution by selection. *Journal of Theoretical Biology*, **110**, 155–172.
- Cheverud, J. M. (1995). Morphological integration in the saddle-back tamarin (*Saguinus fuscicollis*) cranium. *American Naturalist*, **145**, 63–89.
- Ciampaglio, C. N. (2002). Determining the role that ecological and developmental constraints play in controlling disparity: examples from the crinoid and blastozoan fossil record. *Evolution & Development*, **4**, 170–188.
- Ciampaglio, C. N., Kemp, M. and McShea, D. W. (2001). Detecting changes in morphospace occupation patterns in the fossil record: characterization and analysis of measures of disparity. *Paleobiology*, **27**, 695–715.
- Colless, D. H. (1980). Congruence between morphometric and allozyme data for *Menidia* species: a reappraisal. *Systematic Zoology*, **29**, 288–299.
- Cope, E. D. (1887). *The Origin of the Fittest*. McMillan.
- Corruccini, R. S. (1977). Correlation properties of morphometric ratios. *Systematic Zoology*, **26**, 211–214.
- Corti, M. C., Fadda, C., Simson, S. and Nevo, E. (1996). Size and shape variation in the mandible of the fossorial rodent *Spalax ehrenbergi*. In *Advances in Morphometrics* (L. F. Marcus, M. Corti, A. Loy et al., eds) pp. 303–320. Plenum Press.
- Dodson, P. (1975). Relative growth in two sympatric species of *Sceloporus*. *American Midland Naturalist*, **94**, 421–450.
- Dodson, P. (1978). On the use of ratios in growth studies. *Systematic Zoology*, **27**, 62–67.
- Dryden, I. L. and Mardia, K. V. (1998). *Statistical Shape Analysis*. John Wiley & Sons.
- Dryden, I. L. and Walker, G. (1999). Highly resistant regression and object matching. *Biometrics*, **55**, 820–825.
- Edgington, E. S. (1995). *Randomization Tests*. Marcel Dekker.
- Efron, B. (1979). Computers and the theory of statistics, thinking the unthinkable. *Society for Industrial and Applied Mathematics Review*, **21**, 460–480.
- Efron, B. (1987). Better bootstrap confidence-intervals. *Journal of the American Statistical Association*, **82**, 171–185.
- Efron, B. (1992). Jackknife-after-bootstrap standard errors and influence functions. *Journal of the Royal Statistical Society, Series B: Methodological*, **54**, 83–127.
- Efron, B. and Tibshirani, R. J. (1993). *An Introduction to the Bootstrap*. Chapman & Hall.
- Fadda, C., Faggiani, F. and Corti, M. (1997). A portable device for the three-dimensional landmark collection of skeletal elements of small mammals. *Mammalia*, **61**, 622–627.
- Falsetti, A. B. and Cole, T. M. (1992). Relative growth of the postcranial skeleton in callitrichines. *Journal of Human Evolution*, **23**, 79–92.

## BIBLIOGRAPHY

431

- Farris, J. S. (1990). Phenetics in camouflage. *Cladistics*, **6**, 91–100.
- Feller, W. (1968). *An Introduction to Probability Theory and its Applications*. John Wiley & Sons.
- Felsenstein, J. (2002). Quantitative characters, phylogenies, and morphometrics. In *Morphology, Shape and Phylogeny* (N. MacLeod and P. L. Forey, eds) pp. 27–44. Taylor & Francis.
- Ferson, S., Rohlf, F. J. and Koehn, R. K. (1985). Measuring shape variation of two-dimensional outlines. *Systematic Zoology*, **34**, 59–68.
- Fink, W. L. (1993). Revision of the piranha genus *Pygocentrus* (Teleostei, Characiformes). *Copeia*, **1993**, 665–687.
- Fink, W. L. and Zelditch, M. L. (1995). Phylogenetic analysis of ontogenetic shape transformations: a reassessment of the piranha genus *Pygocentrus* (Teleostei). *Systematic Biology*, **44**, 343–360.
- Flury, B. (1988). *Common Principal Components and Related Multivariate Methods*. John Wiley.
- Foote, M. (1986). Developmental buffering as a mechanism for stasis. *Evolution*, **42**, 396–399.
- Foote, M. (1990). Nearest-neighbor analysis of trilobite morphospace. *Systematic Zoology*, **39**, 371–382.
- Foote, M. (1992). Paleozoic record of morphological diversity in blastozoan echinoderms. *Proceedings of the National Academy of Sciences of the United States of America*, **89**, 7325–7329.
- Foote, M. (1993a). Contributions of individual taxa to overall morphological disparity. *Paleobiology*, **19**, 403–419.
- Foote, M. (1993b). Discordance and concordance between morphological and taxonomic diversity. *Paleobiology*, **19**, 185–204.
- Foote, M. (1994). Morphological disparity in Ordovician–Devonian crinoids and the early saturation of morphological space. *Paleobiology*, **20**, 320–344.
- Foote, M. (1997). The evolution of morphological diversity. *Annual Reviews of Ecology and Systematics*, **28**, 129–152.
- Foote, M. and Gould, S. J. (1992). Cambrian and Recent morphological disparity. *Science*, **258**, 816.
- Fuiman, L. A. (1983). Growth gradients in fish larvae. *Journal of Fish Biology*, **23**, 117–123.
- Galis, F. and Metz, J. A. (2001). Testing the vulnerability of the phylotypic stage: on modularity and evolutionary conservatism. *Journal of Experimental Zoology (Molecular and Developmental Evolution)*, **291**, 195–204.
- Galis, F., van Alphen, J. M. and Metz, J. A. (2001). Why five fingers? Evolutionary constraints on digit numbers. *Trends in Ecology and Evolution*, **16**, 637–646.
- Gavrillets, S. (1999). Dynamics of clade diversification on the morphological hypercube. *Proceedings of the Royal Society of London, Series B*, **266**, 817–824.
- Gift, N. and Stevens, P. F. (1997). Vagaries in the delimitation of character states in quantitative variation – an experimental study. *Systematic Biology*, **46**, 112–125.
- Goldman, N. (1988). Methods for discrete coding of morphological characters for numerical analysis. *Cladistics*, **4**, 59–71.
- Good, P. (1994). *Permutation Tests: A Practical Guide to Resampling Methods for Testing Hypotheses*. Springer-Verlag.
- Gould, S. J. (1977). *Ontogeny and Phylogeny*. Harvard University Press.
- Gould, S. J. (1984). Morphological channeling by structural constraint: convergence in styles of dwarfing and gigantism in *Cerion*, with a description of two new fossil species and a report on the discovery of the largest *Cerion*. *Paleobiology*, **10**, 172–194.
- Gould, S. J. and Garwood, R. A. (1969). Levels of integration in mammalian dentitions: an analysis of correlations in *Nesophantes micrus* (Insectivora) and *Oryzomys couesi* (Rodentia). *Evolution*, **23**, 276–300.
- Green, W. D. K. (1996). The thin-plate spline and images with curving features. In *Proceedings in Image Fusion and Shape Variability Techniques* (K. V. Mardia, C. A. Gill and I. L. Dryden, eds) pp. 79–87. Leeds University Press.
- Hall, B. K. (1992). *Evolutionary Developmental Biology*. Chapman & Hall.

- Hills, M. (1978). On ratios – a response to Atchley, Gaskins and Anderson. *Systematic Zoology*, **27**, 61–62.
- Hingst-Zaher, E., Marcus, L. F. and Cerqueria, R. (2000). Application of geometric morphometrics to the study of postnatal size and shape changes in the skull of *Callomys expulsus*. *Hystrix*, **11**, 99–113.
- Hoeffding, W. (1952). The large-sample power of tests based on permutation of observations. *Annals of Mathematical Statistics*, **23**, 169–192.
- Houle, D., Mezey, J. and Galpern, P. (2002). Interpretation of the results of common principal components analysis. *Evolution*, **56**, 433–440.
- Hulsey, C. D. and Wainwright, P. C. (2002). Projecting mechanics into morphospace: disparity in the feeding mechanics of labrid fishes. *Proceedings of the Royal Society of London, Series B*, **269**, 317–326.
- Huxley, J. S. (1932). *Problems of Relative Growth*. MacVeagh.
- Jackson, D. A. and Somers, K. M. (1989). Are probability estimates from the permutation models of Mantel's test stable? *Canadian Journal of Zoology*, **67**, 766–779.
- Jolicoeur, P. (1963). The multivariate generalization of the allometry equation. *Biometrics*, **19**, 497–499.
- Jöreskog, K. G. and Wold, H. (1982). *Systems Under Indirect Observation: Causality–Structure–Prediction*. North Holland Publishing Company.
- Katz, J. M. (1980). Allometry formula: a cellular model. *Growth*, **44**, 89–96.
- Kendall, D. (1977). The diffusion of shape. *Advances in Applied Probability*, **9**, 428–430.
- Kendall, D. G. and Kendall, W. S. (1980). Alignments in two-dimensional random sets of points. *Advances in Applied Probability*, **12**, 380–424.
- Kim, K., Sheets, H. D., Haney, R. A. and Mitchell, C. E. (2002). Morphometric analysis of ontogeny and allometry of the Middle Ordovician trilobite, *Triarthrus becki*. *Paleobiology*, **28**, 364–377.
- Kimmel, C. B., Ballard, W. W., Kimmel, S. R. et al. (1995). Stages of embryonic development of the zebrafish. *Developmental Dynamics*, **203**, 253–310.
- Kingsolver, J. G. and Wiernasz, D. C. (1991). Development, function, and the quantitative genetics of wing melanin pattern in *Pieris* butterflies. *Evolution*, **45**, 1480–1492.
- Klingenberg, C. P. (1998). Heterochrony and allometry: the analysis of evolutionary change in ontogeny. *Biological Reviews*, **73**, 79–123.
- Klingenberg, C. P. and Ekau, W. (1996). A combined morphometric and phylogenetic analysis of an ecomorphological trend: Pelagization in Antarctic fishes (Perciformes: Nototheniidae). *Biological Journal of the Linnean Society*, **59**, 143–177.
- Klingenberg, C. P. and Froese, R. (1991). A multivariate comparison of allometric growth patterns. *Systematic Zoology*, **40**, 410–419.
- Klingenberg, C. P., Badayaev, A. V., Sowry, S. M. and Beckwith, N. J. (2001). Inferring developmental modularity from morphological integration: analysis of individual variation and asymmetry in bumblebee wings. *American Naturalist*, **157**, 11–23.
- Kluge, A. G. and Kerfoot, C. (1973). The predictability and regularity of character divergence. *American Naturalist*, **107**, 426–464.
- Lagler, K. F., Bardach, J. E. and Miller, R. R. (1962). *Ichthyology*. John Wiley & Sons.
- Laird, A. K. (1965). Dynamics of relative growth. *Growth*, **29**, 249–263.
- Laird, A. K., Barton, A. D. and Tyler, S. A. (1968). Growth and time: an interpretation of allometry. *Growth*, **32**, 347–354.
- Lande, R. (1979). Quantitative genetic analysis of multivariate evolution, applied to brain:body size allometry. *Evolution*, **33**, 402–416.
- Lande, R. (1980). The genetic covariance between characters maintained by pleiotropic mutations. *Genetics*, **94**, 314–334.
- Lande, R. and Arnold, S. J. (1983). The measurement of selection on correlated characters. *Evolution*, **37**, 1210–1226.

## BIBLIOGRAPHY

433

- Liebner, D. L. and Sheets, H. D. (2001). Superposer, available on the IMP website [www.canisius.edu/~sheets/morphsoft.html](http://www.canisius.edu/~sheets/morphsoft.html)
- Lohman, G. P. and Schweitzer, P. N. (1990). On eigenshape analysis. In *Proceedings of the Michigan Morphometrics Workshop* (F. J. Rohlf and F. L. Bookstein, eds) pp. 147–166. University of Michigan Museum of Zoology, Special Publication No. 2.
- Lowe, A. A., Özbeck, M. M., Miyamoto, K. and Fleetham, J. A. (1997). Cephalometric and demographic characteristics of sleep apnea: an evaluation with partial least squares analysis. *The Angle Orthodontist*, **67**, 143–154.
- Lundrigan, B. (1996). Morphology of horns and fighting behavior in the family Bovidae. *Journal of Mammalogy*, **77**, 462–475.
- MacLeod, N. and Rose, K. D. (1993). Inferring locomotor behavior in Paleogene mammals via eigenshape analysis. *American Journal of Science*, **293A**, 300–355.
- Manly, B. F. (1997). *Randomization, Bootstrap and Monte Carlo Methods in Biology*. Chapman & Hall.
- Marcus, L. F., Hingst-Zaher, E. and Zaher, H. (2000). Applications of landmark morphometrics to skulls representing the orders of living mammals. *Hystrix* (n.s.), **11**, 24–48.
- Marroig, G. and Cheverud, J. M. (2001). A comparison of phenotypic variation and covariation patterns and the role of phylogeny, ecology and ontogeny during cranial evolution of New World monkeys. *Evolution*, **55**, 2576–2600.
- Maynard Smith, J. M., Burian, R., Kauffman, S. et al. (1984). Developmental constraints and evolution. *Quarterly Review of Biology*, **60**, 265–287.
- McKinney, M. L. and McNamara, K. J. (1991). *Heterochrony: The Evolution of Ontogeny*. Plenum Press.
- Miller, A. I. and Foote, M. (1996). Calibrating the Ordovician radiation of marine life: implications for Phanerozoic diversity trends. *Paleobiology*, **22**, 304–309.
- Morrison, D. F. (1990). *Multivariate Statistical Methods*. McGraw-Hill.
- Myers, P., Lundrigan, B. L., Gillespie, B. W. and Zelditch, M. L. (1996). Phenotypic plasticity in skull and dental morphology in the Prairie Deer Mouse (*Peromyscus maniculatus bairdii*). *Journal of Morphology*, **229**, 229–237.
- Olson, E. C. and Miller, R. L. (1958). *Morphological Integration*. University of Chicago Press.
- Oxnard, C. E. (1968). The architecture of the shoulder in some mammals. *Journal of Morphology*, **126**, 249–290.
- Phillips, P. C. and Arnold, S. J. (1999). Hierarchical comparison of genetic variance–covariance matrices. I. Using the Flury hierarchy. *Evolution*, **53**, 1506–1515.
- Polly, P. D. (200x). Geography and sample size in P matrix evolution: molar shape change in island populations of *Sorex araneus*. *Journal of Evolutionary Biology*.
- Press, W. H., Flannery, B. P., Teukolsky, S. A. and Vetterling, W. T. (1998). *Numerical Recipes in C: The Art of Scientific Computing*. Cambridge University Press.
- Quenouille, M. (1949). Approximate tests of correlation in time series. *Journal of the Royal Statistical Society B*, **11**, 18–14.
- Raff, R. A. (1992). Direct-developing sea urchins and the evolutionary reorganization of early development. *BioEssays*, **14**, 211–218.
- Rao, C. R. (1973). *Linear Statistical Inference and its Applications*. John Wiley & Sons.
- Richardson, M. K., Hanken, J., Gooneratne, M. L. et al. (1997). There is no highly conserved embryonic stage in the vertebrates: implications for current theories of evolution and development. *Anatomy and Embryology*, **196**, 91–106.
- Robinson, J. (1973). Large-sample power of permutation tests for randomization models. *Annals of Statistics*, **1**, 291–296.
- Rohlf, F. J. (1990). Rotational fit (Procrustes) methods. In *Proceedings of the Michigan Morphometrics Workshop* (F. J. Rohlf and F. L. Bookstein, eds) pp. 227–236. University of Michigan Museum of Zoology, Special Publication No. 2.

- Rohlf, F. J. (1998). On applications of geometric morphometrics to studies of ontogeny and phylogeny. *Systematic Biology*, **47**, 147–158.
- Rohlf, F. J. (2000). On the use of shape spaces to compare morphometric methods. *Hystrix* (n.s.), **11**, 8–24.
- Rohlf, F. J. (2002). Geometric morphometrics and phylogeny. In *Morphology, Shape and Phylogeny* (N. MacLeod and P. L. Forey, eds) pp. 175–193. Taylor & Francis.
- Rohlf, F. J. and Archie, J. W. (1984). A comparison of Fourier methods for the description of wing shape in mosquitoes (Diptera: Culicidae). *Systematic Zoology*, **33**, 302–317.
- Rohlf, F. J. and Bookstein, F. L. (2003). Computing the uniform component of shape variation. *Systematic Biology*, **52**, 66–69.
- Rohlf, F. J. and Corti, M. (2000). Use of two-block partial least squares to study covariation in shape. *Systematic Biology*, **49**, 740–753.
- Rohlf, F. J. and Slice, D. E. (1990). Extensions of the Procrustes method for the optimal superimposition of landmarks. *Systematic Zoology*, **39**, 40–59.
- Rohlf, F. J., Gilmartin, A. J. and Hart, G. (1983). The Kluge–Kerfoot phenomenon: a statistical artifact? *Evolution*, **37**, 180–202.
- Romano, J. P. (1989). Bootstrap and randomization tests of some non-parametric hypotheses. *Annals of Statistics*, **17**, 141–159.
- Roth, V. L. (1993). On three-dimensional morphometrics, and on the identification of landmark points. In *Contributions to Morphometrics* (L. F. Marcus, E. Bello and A. García-Valdecasas, eds), pp. 41–62. Museo Nacional de Ciencias Naturales, Madrid.
- Roy, K. and Foote, M. (1997). Morphological approaches to measuring biodiversity. *Trends in Ecology & Evolution*, **12**, 277–281.
- Rüber, L. and Adams, D. C. (2001). Evolutionary convergence of body shape and trophic morphology in cichlids of Lake Tanganyika. *Journal of Evolutionary Biology*, **14**, 325–332.
- Sampson, P. D., Streissguth, A. P., Barr, H. M. and Bookstein, F. L. (1989). Neurobehavioral effects of prenatal alcohol: Part II. Partial least squares analysis. *Neurotoxicology and Teratology*, **11**, 477–491.
- Sampson, P. D., Bookstein, F. L., Sheehan, F. H. and Bolson, E. L. (1996). Eigenshape analysis of left ventricular outlines from contrast ventriculograms. In *Advances in Morphometrics* (L. F. Marcus, M. Corti, A. Loy et al., eds) pp. 211–233. Plenum.
- Sander, K. (1983). The evolution of patterning mechanisms: gleanings from insect embryogenesis and spermatogenesis. In *Development and Evolution* (B. C. Goodwin, N. Holder and C. C. Wylie, eds) pp. 137–159. Cambridge University Press.
- Schaefer, S. A. and Lauder, G. V. (1996). Testing historical hypotheses of morphological change: biomechanical decoupling in loricariid catfishes. *Evolution*, **50**, 1661–1675.
- Schluter, D. (1996). Adaptive radiation along genetic lines of least resistance. *Evolution*, **50**, 1766–1774.
- Seidl, F. (1960). Körpergrundgestalt und Keimstruktur. Eine Erörterung über die Grundlagen der vergleichenden und experimentellen Embryologie und deren Gültigkeit bei phylogenetischen Überlegungen. *Zoologische Anzeiger*, **164**, 245–305.
- Shea, B. T. (1992). Ontogenetic scaling of skeletal proportions in the talapoin monkey. *Journal of Human Evolution*, **23**, 283–307.
- Shea, B. T. (2002). Are some heterochronic transformations likelier than others? In *Human Evolution Through Developmental Change* (N. Minugh-Purvis and K. J. McNamara, eds) pp. 79–101. Johns Hopkins Press.
- Sheets, H. D. and Mitchell, C. E. (2001). Why the null matters: statistical tests, random walks and evolution. *Genetica*, **112**, 105–125.
- Siegel, A. F. and Benson, R. H. (1982). A robust comparison of biological shapes. *Biometrics*, **38**, 341–350.
- Simon, C. (1983). A new coding procedure for morphometric data with an example from periodical cicada wing veins. In *Numerical Taxonomy* (J. Felsenstein, ed.) pp. 378–382. Springer-Verlag.

## BIBLIOGRAPHY

435

- Slack, J. M., Holland, P. W. and Graham, C. F. (1993). The zootype and the phylotypic stage. *Nature*, **361**, 490–492.
- Slice, D. E. (2001). Landmark coordinates aligned by Procrustes analysis do not lie in Kendall's shape space. *Systematic Biology*, **50**, 141–149.
- Slice, D. E., Bookstein, F. L., Marcus, L. F. and Rohlf, F. J. (1996). Appendix I: A glossary for geometric morphometrics. In *Advances in Morphometrics* (L. F. Marcus, M. Corti, A. Loy et al., eds) pp. 531–551. Plenum Press.
- Small, C. G. (1996). *The Statistical Theory of Shape*. Springer.
- Smith, L. H. and Lieberman, B. S. (1999). Disparity and constraint in olenelloid trilobites and the Cambrian radiation. *Paleobiology*, **25**, 459–470.
- Snedecor, G. W. and Cochran, W. G. (1967). *Statistical Methods*. Iowa State University Press.
- Sokal, R. R. (1976). The Kluge–Kerfoot phenomenon re-examined. *American Naturalist*, **110**, 1077–1091.
- Sokal, R. R. and Rohlf, F. J. (1995). *Biometry: The Principals and Practice of Statistics in Biological Research*, 3rd edn. Freeman.
- Spencer, M. A. and Spencer, G. S. (1995). Video-based three-dimensional morphometrics. *American Journal of Physical Anthropology*, **96**, 443–453.
- Stein, B. R. (1981). Comparative limb myology of two opossums, *Didelphis* and *Chironectes*. *Journal of Morphology*, **169**, 113–140.
- Steppan, S. J. (1997). Phylogenetic analysis of phenotypic covariance structure. I. Contrasting results from matrix correlation and common principal component analysis. *Evolution*, **51**, 571–586.
- Strauss, D. and Sadler, P. M. (1989). Classical confidence intervals and Bayesian probability estimates for ends of local taxon ranges. *Mathematical Geology*, **21**, 411–427.
- Strauss, R. E. (1984). Allometry and functional feeding morphology in haplochromine cichlids. In *Evolution of Fish Species Flocks* (A. A. Echelle and I. Kornfield, eds) pp. 217–229. University of Maine.
- Strauss, R. E. and Bookstein, F. L. (1982). The truss – body form reconstructions in morphometrics. *Systematic Zoology*, **31**, 113–135.
- Strauss, R. E. and Fuiman, L. A. (1985). Quantitative comparisons of body form and allometry in larval and adult Pacific sculpins (Teleostei: Cottidae). *Canadian Journal of Zoology*, **63**, 1582–1589.
- Streissguth, A. P., Bookstein, F. L., Sampson, P. D. and Barr, H. M. (1993). *The Enduring Effects of Prenatal Alcohol Exposure on Child Development: Birth Through Seven Years, A Partial Least Squares Solution*. University of Michigan Press.
- Swiderski, D. L. (1993). Morphological evolution of the scapula in tree squirrels, chipmunks, and ground squirrels (Sciuridae): an analysis using thin-plate splines. *Evolution*, **47**, 1854–1873.
- Swiderski, D. L., Zelditch, M. L. and Fink, W. L. (1998). Why morphometrics isn't special: coding quantitative data for phylogenetic analysis. *Systematic Biology*, **47**, 508–519.
- Taylor, M. E. (1974). The functional anatomy of the forelimb of some African Viverridae (Carnivora). *Journal of Morphology*, **143**, 307–336.
- Thiele, K. (1993). The holy grail of the perfect character: the cladistic treatment of morphometric data. *Cladistics*, **9**, 275–304.
- Thompson, D'Arcy W. (1942). *On Growth and Form: A New Edition*. Cambridge University Press. (Reprinted in 1992 as *On Growth and Form: The Complete Revised Edition*, Dover Publications.)
- Tukey, J. W. (1958). Bias and confidence in not quite large samples (Abstract). *Annals of Mathematical Statistics*, **29**, 614.
- van Snik, G. M. J., van den Boogaart, J. G. M. and Osse, W. M. (1997). Larval growth patterns in *Cyprinus carpio* and *Clarias gariepinus* with attention to the finfold. *Journal of Fish Biology*, **50**, 1339–1352.
- Van Valen, L. (1962). Developmental gradients in the dentition of *Peromyscus*. *Evolution*, **16**, 272–277.
- Van Valen, L. (1970). An analysis of developmental fields. *Developmental Biology*, **23**, 456–477.

- Voss, R. S. and Marcus, L. F. (1992). Morphological evolution in muroid rodents.2. Cranio-metric factor divergence in 7 neotropical genera, with experimental results from *Zygodontomys*. *Evolution*, **46**, 1918–1934.
- Wagner, G. P. (1988). The influence of variation and of developmental constraints on the rate of multivariate phenotypic evolution. *Journal of Evolutionary Biology*, **1**, 45–66.
- Wagner, G. P. and Altenberg, L. (1996). Complex adaptations and the evolution of evolvability. *Evolution*, **50**, 967–976.
- Wagner, P. J. (1995). Testing evolutionary constraint hypotheses with early Paleozoic gastropods. *Paleobiology*, **21**, 248–272.
- Wagner, P. J. (1997). Patterns of morphologic diversification among the Rostroconchia. *Paleobiology*, **23**, 115–150.
- Walker, J. A. (2000). Ability of geometric morphometric methods to estimate a known covariance matrix. *Systematic Biology*, **49**, 686–696.
- Wayne, R. K. (1986). Cranial morphology of domestic and wild canids: the influence of development on morphological change. *Evolution*, **40**, 243–261.
- Webster, M., Sheets, H. D. and Hughes, N. C. (2001). Allometric patterning in trilobite ontogeny: testing for heterochrony in *Nephrolenellus*. In *Beyond Heterochrony: The Evolution of Development* (M. L. Zelditch, ed.) pp. 105–142. John Wiley & Sons.
- White, J. F. and Gould, S. J. (1965). Interpretation of the coefficients in the allometric equation. *American Naturalist*, **99**, 5–18.
- Wills, M. A. (2001). Morphological disparity: a primer. In *Fossils, Phylogeny, and Form: An Analytical Approach* (J. M. Adrain, G. D. Edgecombe and B. S. Lieberman, eds) pp. 55–144. Kluwer Academic/Plenum Publishers.
- Wills, M. A., Briggs, D. E. G. and Fortey, R. A. (1994). Disparity as an evolutionary index – a comparison of Cambrian and recent arthropods. *Paleobiology*, **20**, 93–130.
- Zelditch, M. L. (1988). Ontogenetic variation in patterns of phenotypic integration in the laboratory rat. *Evolution*, **42**, 28–41.
- Zelditch, M. L. and Carmichael, A. C. (1989). Ontogenetic variation in patterns of developmental and functional integration in skulls of *Sigmodon fulviventer*. *Evolution*, **43**, 814–824.
- Zelditch, M. L., Bookstein, F. L. and Lundrigan, B. L. (1992). Ontogeny of integrated skull growth in the cotton rat *Sigmodon fulviventer*. *Evolution*, **46**, 1164–1180.
- Zelditch, M. L., Bookstein, F. L. and Lundrigan, B. L. (1993). The ontogenetic complexity of developmental constraints. *Journal of Evolutionary Biology*, **6**, 121–141.
- Zelditch, M. L., Fink, W. L. and Swiderski, D. L. (1995). Morphometrics, homology and phylogenetics: quantified characters as synapomorphies. *Systematic Biology*, **44**, 179–189.
- Zelditch, M. L., Fink, W. L., Swiderski, D. L. and Lundrigan, B. L. (1998). On applications of geometric morphometrics to studies of ontogeny and phylogeny: a reply to Rohlf. *Systematic Biology*, **47**, 159–167.
- Zelditch, M. L., Sheets, H. D. and Fink, W. L. (2000). Spatiotemporal reorganization of growth rates in the evolution of ontogeny. *Evolution*, **54**, 1363–1371.
- Zelditch, M. L., Sheets, H. D. and Fink, W. L. (2001). The spatial complexity and evolutionary dynamics of growth. In *Beyond Heterochrony: The Evolution of Development* (M. L. Zelditch, ed.) pp. 145–194. John Wiley & Sons.
- Zelditch, M. L., Sheets, H. D. and Fink, W. L. (2003b). The ontogenetic dynamics of shape disparity. *Paleobiology*, **29**, 139–156.
- Zelditch, M. L., Lundrigan, B. L., Sheets, H. D. and Garland, T. Jr (2003b). Do precocial mammals develop at a faster rate? A comparison of rates of skull development in *Sigmodon fulviventer* and *Mus musculus domesticus*. *Journal of Evolutionary Biology*, **16**, 708–720.