

COMMENT

No rattlesnakes in the rainforests: reply to Gosling and Bush

WOLFGANG WÜSTER,* JULIA E. FERGUSON,* J. ADRIAN QUIJADA-MASCAREÑAS,*
CATHARINE E. POOK,* MARIA DA GRAÇA SALOMÃO† and ROGER S. THORPE*

*School of Biological Sciences, University of Wales, Bangor LL57 2UW, Wales, UK, †Laboratório de Herpetologia, Instituto Butantan, Avenida Vital Brazil 1500, 05503-900 São Paulo — SP, Brazil

Gosling & Bush (2005) criticize our interpretation (Wüster *et al.* 2005) of a phylogeographic analysis of the Neotropical rattlesnake (*Crotalus durissus*) as supporting a hypothesis of Pleistocene fragmentation of the Amazonian rainforests. We thank these two authors for the opportunity to further discuss issues concerning the ecological and biogeographical history of the Amazon Basin.

Gosling and Bush quite rightly comment on the diversity of the ecosystems that are often grouped together as 'Amazonian rainforest', and also note the dynamic changes of community composition throughout the Pleistocene. They go on to comment that these two factors make any extrapolation from present-day distribution patterns difficult, and question the use of *C. durissus* to determine the possible past extent of rainforest fragmentation. We reject these doubts, and maintain that the presence of *C. durissus* on both sides of the Amazon Basin does represent evidence of profound changes in the distribution of rainforests in the Pleistocene and cannot be explained by relatively minor changes in rainforest community composition.

Despite the dynamic nature of community composition in Neotropical forests, a typical rainforest physiognomy can be defined and can be traced back through the vegetational history of South America to at least the Eocene (Burnham & Johnson 2004). Moreover, the many different identifiable Amazonian forest types all have one factor in common: *C. durissus* does not occur in them. The few available records of *C. durissus* in forest zones (e.g. Beebe 1946) are all of occasional specimens from ecotonal zones close to established populations in open formations. Kartabo (6°23'N, 58°41'W), the site of Beebe's records, is situated at the edge of the documented distribution of *C. durissus* in Guyana (Campbell & Lamar 2004). Numerous intensive herpetofaunal surveys in many parts of the Amazon and adjacent moist forests, including many different forest types, have consistently failed to locate *C. durissus* in rainforest areas (e.g. Duellman 1978; Dixon &

Soini 1986; Cunha & Nascimento 1993; Martins & Oliveira 1998), including in other parts of Guyana (Donnelly *et al.* 2005), and no established moist forest populations are known (Campbell & Lamar 2004). On the other hand, *C. durissus* has proven quick to occupy anthropogenically deforested areas in tropical South America (Sazima & Haddad 1992; Melgarejo & Aguiar 1995; Marques *et al.* 1998; Antunes 2003). Altogether, this suggests that *C. durissus* is a species that is unable to persist in a wide range of moist forest types.

We reject the suggestion that *C. durissus* may have occupied rainforest formations in the past as entirely unsupported. Apart from the absence of forest-dwelling populations at present, we also note that no other species of *Crotalus* occupies tropical rainforests (Campbell & Lamar 2004). Indeed, the closest relatives of *C. durissus*, such as *Crotalus molossus* and *Crotalus basiliscus*, occupy highly seasonal or xeric formations (Campbell & Lamar 2004), and the ancestral habitat for rattlesnakes is most likely to have been upland pine-oak forest (Place & Abramson 2004), a much more open formation than any Amazonian rainforest. This makes an assumption of Pleistocene rainforest occupancy by *C. durissus* highly unparsimonious.

Gosling & Bush (2005) suggest several alternative scenarios for the present-day distribution of *C. durissus*. They back these scenarios up with late Pleistocene palynological information and climatic models for the Last Glacial Maximum. The suggestion that the forests of eastern Amazonia, which surround the isolated Santarém population of *C. durissus*, may have been particularly susceptible to opening up as a result of reduced moisture during the Pleistocene is precisely the kind of scenario our data support: it clearly implies the isolation of the Guyanan rainforests from those of the western Amazon Basin, i.e. rainforest fragmentation, perhaps as illustrated by Hooghiemstra & van der Hammen (1998; Fig. 3).

However, in any case, we re-emphasize that our data on *C. durissus* suggest that forest fragmentation occurred in the Middle Pleistocene, and thus provide no information on late Pleistocene events. Published palynological data have increasingly been interpreted as contradicting extensive

rainforest fragmentation in the Late Pleistocene (Colinvaux *et al.* 1996, 2001; Haberle & Maslin 1999; Kastner & Goñi 2003; Bush *et al.* 2004), but pollen records from the Early and Middle Pleistocene are lacking. There is therefore no palynological evidence to reject a hypothesis of rainforest fragmentation in the Middle or Early Pleistocene.

The precise nature and extent of this fragmentation remain to be ascertained: our data do not necessarily call for the same extent of rainforest fragmentation as the early formulations of the refugia hypothesis (Haffer 1969), and neither do they predict to what extent any intervening vegetation would have consisted of deciduous forests (Prado & Gibbs 1993), tree savannahs (e.g. Rossetti *et al.* 2004), savannah–forest mosaics (Webb & Rancy 1996) or true savannahs (Haffer 1969). All of these provide suitable habitat for *C. durissus*, and all are inconsistent with the view that the Amazon Basin remained cloaked in rainforest throughout the Pleistocene (e.g. Colinvaux *et al.* 2000, 2001; Bush *et al.* 2004).

Lastly, the alternative scenarios suggested by Gosling & Bush (2005) to account for the present-day distribution of *C. durissus* are to a large extent just-so stories based on speculation rather than data. In contrast, our hypothesis of mid-Pleistocene forest fragmentation generated on the basis of the *C. durissus* data makes clear, testable predictions of patterns that we may expect to see in other nonrainforest organisms distributed on both sides of the Amazon Basin. The very similar findings of Eberhard & Bermingham (2004) in *Amazona ochrocephala* and the existence of multiple open-formation plant species distributed on different sides of the Amazonian evergreen forests, including both north and south of the Amazon Basin (Prado & Gibbs 1993; Pennington *et al.* 2000, 2004), encourage us in our belief that intraspecific phylogeographic studies of nonrainforest organisms, complementing the interspecific approach adopted by Pennington *et al.* (2004), have much to contribute to the debate about the history of the New World tropics. We thus hope that our study of the phylogeography of *C. durissus* will stimulate further studies along similar lines to test for the existence of common patterns that may reveal common causes.

Received 22 May 2005; revision accepted 2 June 2005

References

- Antunes AP (2003) Colonização por serpentes em área alterada da Serra do Mar, Município de Pilar do Sul-SP. In: *Anais do VI Congresso de Ecologia do Brasil, Fortaleza*, pp. 179–180. Fortaleza, Brazil.
- Beebe W (1946) Field notes on the snakes of Kartabo, British Guiana and Caripito, Venezuela. *Zoologica*, **31**, 11–52.
- Burnham RJ, Johnson KR (2004) South American palaeobotany and the origins of neotropical rainforests. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, **359**, 1595–1610.
- Bush MB, de Oliveira PE, Colinvaux PA *et al.* (2004) Amazonian paleoecological histories: one hill, three watersheds. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **214**, 359–393.
- Campbell JA, Lamar WW (2004) *The Venomous Reptiles of the Western Hemisphere*. Comstock Publishing Associates, Ithaca and London.
- Colinvaux PA, De Oliveira PE, Moreno JE, Miller MC, Bush MB (1996) A long pollen record from lowland Amazonia: forest and cooling in glacial times. *Science*, **274**, 85–87.
- Colinvaux PA, De Oliveira PE, Bush MB (2000) Amazonian and Neotropical plant communities on glacial time-scales: the failure of the aridity and refuge hypotheses. *Quaternary Science Reviews*, **19**, 141–169.
- Colinvaux PA, Irion G, Räsänen ME, Bush MB, Nunes de Mello JAS (2001) A paradigm to be discarded: geological and paleoecological data falsify the Haffer & Prance refuge hypothesis of Amazonian speciation. *Amazoniana*, **16**, 609–646.
- Cunha OR, Nascimento FP (1993) Ofídios da Amazônia. As cobras da região leste do Pará. *Boletim do Museu Paraense Emílio Goeldi*, **9**, 1–191.
- Dixon JR, Soini P (1986) *The Reptiles of the Upper Amazon Basin, Iquitos Region, Peru*. Milwaukee Public Museum, Milwaukee, Wisconsin.
- Donnelly MA, Chen MH, Watkins GG (2005) Sampling amphibians and reptiles in the Iwokrama Forest ecosystem. *Proceedings of the Academy of Natural Sciences of Philadelphia*, **154**, 55–69.
- Duellman WE (1978) The biology of an equatorial herpetofauna in Amazonian Ecuador. *Miscellaneous Publications of the Museum of Natural Science, University of Kansas*, **65**, 1–352.
- Eberhard JR, Bermingham E (2004) Phylogeny and biogeography of the *Amazona ochrocephala* (Aves: Psittacidae) complex. *Auk*, **121**, 318–332.
- Gosling WD, Bush MB (2005) A biogeographic comment on: Wüster, W *et al.* (2005) Tracing an invasion: landbridges, refugia, and the phylogeography of the Neotropical rattlesnake (Serpentes: Viperidae: *Crotalus durissus*). *Molecular Ecology*, in press.
- Haberle SG, Maslin MA (1999) Late Quaternary vegetation and climate change in the Amazon Basin based on a 50 000-year pollen record from the Amazon Fan, ODP Site 932. *Quaternary Research*, **51**, 27–38.
- Haffer J (1969) Speciation in Amazonian forest birds. *Science*, **165**, 131–137.
- Hooghiemstra H, van der Hammen T (1998) Neogene and Quaternary development of the Neotropical rain forest: the forest refugia hypothesis and a literature overview. *Earth Science Reviews*, **44**, 147–183.
- Kastner TP, Goñi MA (2003) Constancy in the vegetation of the Amazon Basin during the late Pleistocene: evidence from the organic matter composition of Amazon deep sea fan sediments. *Geology*, **31**, 291–294.
- Marques OAV, Abe AS, Martins M (1998) Estudo diagnóstico da diversidade de répteis do Estado de São Paulo. In: *Biodiversidade do Estado de São Paulo, Brasil, Vol. 6, Vertebrados* (ed. Castro RMC), pp. 27–38. FAPESP, São Paulo.
- Martins M, Oliveira ME (1998) Natural history of snakes in forests of the Manaus Region, Central Amazonia, Brazil. *Herpetological Natural History*, **6**, 78–150.
- Melgarejo AR, Aguiar AS (1995) Poisonous snakes, ecological disturbances and public health. In: *1st International Congress on Envenomations and Their Treatments*, p. 49. Institut Pasteur, Paris.
- Pennington RT, Prado DE, Pendry CA (2000) Neotropical seasonally dry forests and Quaternary vegetation changes. *Journal of Biogeography*, **27**, 261–273.

- Pennington RT, Lavin M, Prado DE *et al.* (2004) Historical climate change and speciation: Neotropical seasonally dry forest plants show patterns of both Tertiary and Quaternary diversification. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, **359**, 515–538.
- Place AJ, Abramson CI (2004) A quantitative analysis of the ancestral area of rattlesnakes. *Journal of Herpetology*, **38**, 151–156.
- Prado DE, Gibbs PE (1993) Patterns of species distributions in the dry seasonal forests of South America. *Annals of the Missouri Botanical Garden*, **80**, 902–927.
- Rossetti DF, Toledo PM, Moraes-Santos HM, Santos AEA (2004) Reconstructing habitats in central Amazonia using megafauna, sedimentology, radiocarbon, and isotope analyses. *Quaternary Research*, **61**, 289–300.
- Sazima I, Haddad CFB (1992) Répteis da Serra do Japi: notas sobre historia natural. In: *História Natural da Serra do Japi. Ecologia e Preservação de uma Área Florestal no Sudeste do Brasil* (ed. Morellato LPC), pp. 212–326. Editora da UNICAMP/FAPESP, Campinas.
- Webb SD, Rancy A (1996) Late Cenozoic evolution of the Neotropical mammal fauna. In: *Evolution and Environment in Tropical America* (eds Jackson JBC, Budd AF, Coates AG), pp. 335–358. University of Chicago Press, Chicago.
- Wüster W, Ferguson JE, Quijada-Mascareñas A, Pook CE, Salomão MG, Thorpe RS (2005) Tracing an invasion: landbridges, refugia, and the phylogeography of the Neotropical rattlesnake (Serpentes: Viperidae: *Crotalus durissus*). *Molecular Ecology*, **14**, 1095–1108.
-
- This study forms part of a wide-ranging collaborative research program into the evolutionary biology, phylogeny and biogeography of venomous snakes and Neotropical biogeography by the authors.
-